

# **Quantitative Evaluations of the Effects of the Seabed Sediments on the Scattering and Propagation of Low to Medium Frequency Acoustic Energy in the Shallow Oceans**

Tokuo Yamamoto

Geoacoustical Laboratory, RSMAS, University of Miami

4600 Rickenbacker Causeway

Miami, FL 33149

phone: (305) 361-4637 fax: (305) 361-4781 email: [tyamamoto@rsmas.miami.edu](mailto:tyamamoto@rsmas.miami.edu)

Award Number: N000140010032

<http://www.rsmas.miami.edu/div/amp/geoacoustics>

Thrust Category: Shallow-Water Acoustics

## **LONG-TERM GOALS**

To quantitatively understand the physics of propagation and scattering of low to medium frequency (20-15000 Hz) acoustic wave in shallow waters and seabed sediments.

## **OBJECTIVES**

The first is to quantitatively understand the effects of seabed scattering mechanisms (volume fluctuation, bottom and sub-bottom roughness) on the acoustic propagation and scattering. This includes the effects of poro-elastic properties of the sediments on the propagation of acoustic waves. The secondary objective is the quantitative understanding of the spatial and temporal fluctuations (amplitude and phase coherency) of acoustic wave fields in shallow water.

## **APPROACH**

The monostatic and bistatic scattering from seabed sediments will be measured using our bilinear array in sandy bottom and clay bottom. At the same time the reciprocal transmission of Gold code acoustic signals at center frequency at 5.5 kHz (1023 cycles, three sine waves per cycle) will be made among three stations separated by 5 to 20 km to study the effect of bottom sediments, spatial and temporal fluctuations within the water column. Inversely, these acoustic data will be inverted to extract sediment properties and water column physical properties of internal waves, tides and fronts. These measurements will be compared with the low frequency scattering and propagation measurements by Dr. Kazi Ohta, of Japan Defense Agency, Dr. Altan Turgut of NRL, and Dr. George Frisk of WHOI.

## **WORK COMPLETED**

Three autonomous acoustic reciprocal transmitters have been built and being tested for the SWAT October 2000 experiment. One of them is also equipped with 32-channel bilinear array for monostatic and bistatic scattering measurements.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>SEP 2000</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2000 to 00-00-2000</b>	
4. TITLE AND SUBTITLE <b>Quantitative Evaluations of the Effects of the Seabed Sediments on the Scattering and Propagation of Low to Medium Frequency Acoustic Energy in the Shallow Oceans</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Geoacoustical Laboratory, RSMAS, University of Miami ,4600 Rickenbacker Causeway ,Miami,,FL,33149</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>4</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## RESULTS

Monostatic scattering and bistatic scattering data at 15 kHz from a sandy bottom and a clay bottom are critically compared. It was learned that at small grazing angle (less than 15 degree ) the bistatic scattering is stronger than the monostatic scattering regardless the sediment type (Day and Yamamoto, 2000; Yamamoto and Sun, 2000). The monostatic scattering at large grazing angles from sediments is mainly from the sediment volume (Rogers and Yamamoto, 1999; Day and Yamamoto, 1999).

The apparent attenuation within the nearsurface sediments at medium to high frequency is due to the Biot mechanism and the scattering from the velocity and density fluctuation within sediment volume according to the velocity and attenuation measurements.(Yamamoto, INPRESS)

## IMPACT/APPLICATIONS

Monostatic scattering measurements from sediments closely follows the anisotropic volume velocity and density fluctuation represented by Yamamoto (1996).

The permeability structure within the near surface sediments was imaged for the first time by crosswell tomography (Yamamoto, I N PRESS).

## TRANSITIONS

The Yamamoto model of anisotropic velocity and density fluctuations is widely used in 6.2 and 6.3 community for scattering modeling.

## RELATED PROJECTS

We collaborate with Dr. Kazi Ohta, of Japan Defense Agency, Dr. Altan Turgut of NRL, and Dr. George Frisk of WHOI.on the research of mutual interest on bottom sediment characterization, scattering and propagation.

## REFERENCES

1. Day, C. M. and Yamamoto, T., "Low graging angle bistatic sea floor scattering on the Florida Atlantic coastal shelf," J. Acoust. Soc. Am., **106 (4)**, pp1744-1754.
2. Rogers, A.K., and Yamamoto, T., "Analysis of high-frequency acoustic scattering data measured in the shallow waters of the Florida Strait, J. Acoust. Soc. Am., **106 (5)**, pp2469-2480.
3. Yamamoto, T, "Imaging the permeability structure within the near-surface sediments by acoustic crosswell tomography," J. Applied Geophysics [IN PRESS].
4. Yamamoto,T. and Sun, H., "Critical comparisions of the measured forward scattering from a clay bottom and a sandy bottom.", Paper #5aUW5, 139<sup>th</sup> Meeting: Acoustical Society of America, Spring Meeting 2000.

## **PUBLICATIONS**

1. Day, C. M. and Yamamoto, T., “Low grazing angle bistatic sea floor scattering on the Florida Atlantic coastal shelf,” J. Acoust. Soc. Am., **106 (4)**, pp1744-1754.
2. Rogers, A.K., and Yamamoto, T., “Analysis of high-frequency acoustic scattering data measured in the shallow waters of the Florida Strait, J. Acoust. Soc. Am., **106 (5)**, pp2469-2480.
3. Yamamoto, T, “Imaging the permeability structure within the near-surface sediments by acoustic crosswell tomography,” J. Applied Geophysics [IN PRESS].
4. Yamamoto, T. and Sun, H., “Critical comparisons of the measured forward scattering from a clay bottom and a sandy bottom.”, Paper #5aUW5, 139<sup>th</sup> Meeting: Acoustical Society of America, Spring Meeting 2000.

## **PATENTS**

Yamamoto, T., #6,061,3000 May 9. 2000. “Method of Imaging the Permeability and Fluid Content Structure within Sediment.”